

Aquilaria malaccensis Lam.

Syn. *A. agallocha* Roxb.

Fam. Thymelaeaceae

Ayurvedic name	Agaru
Unani name	Ood Hindi /Agar
Hindi name	Agar
English name	Aloewood, Eaglewood, Agarwood
Trade name	Agar
Parts used	Fragrant Resinous Wood and Oil



Aquilaria malaccensis

Morphological Characteristics

This plant is a large evergreen tree about 20 meters tall and 1.5–2.4 meters in girth with somewhat straight and fluted bole. Leaves are alternate 0.5-10 cm by 2-5 cm, oblong, lanceolate or elliptic, caudate, acuminate and glabrous with slender nerves. Venation is parallel. Petiole is 0.3-0.5 cm long. It is commercially used as fragrant and in preparation of drugs. The tree contains plenty of oleoresin and has irregular dark patches. The wood burns with a bright flame giving off pleasant smell.

Floral Characteristics

Flowers are white in colour, bisexual, pedicellate, in both axillary and terminal umbellate cymes, shortly peduncled, perianth, campanulate, lobes 5 spreading and densely pilose. Pedicels is 0.5-0.8 cm long, slender. Perianth remains persistent in fruit and 1.3-1.5 cm long, silky densely villous, connate at the base. Stamens are 10, anthers 10 with subsessile disc. Ovary is subsessile, villous and two-celled. Stigma is large, subsessile. Fruit is capsular, 3-5 cm long, obovoid, pericarp coriaceous and densely tomentose. Seeds are ovoid with a long tail.

Distribution

Bengal and North-Eastern States of India namely Assam, Meghalaya, Manipur, Mizoram,



- **Disease and Pest Control:** Attack of *Heortia vitessoides* is observed during May-August. This causes defoliation of whole tree. Application of Thiodan @ 2 ml/lit at 15 days interval during infestation is found to control the pests effectively.

Harvest Management

Agar-wood develops a peculiar, persisting strong odour because of infestation by a fungal identified as *Zeuzera conferta*, it penetrates the hard wood, through wounds, injury or borers. All attempts to induce artificial infestation have failed; it is a natural phenomenon. It develops black patches and stores resinous oil which is separated through distillation of the woody chips. This oil has high value in medicine and perfumery industry.

- **Crop Maturity and Harvesting:** Time of harvesting depends on disease infestation in hard wood. Agar is regarded as a pathological product formed as result of infection. Black patches in the bark indicate occurrence of infection and can be used for harvesting hard wood to commercial use.
- **Post-harvest Management:** Wood chips or chips powdered mechanically without generating heat are soaked in water for 2-3 days and transferred to stainless steel vessel which is part of a distillation unit. The distillation is done for 30-36 hours. Oil and water is collected in a separator and stored. The oil and water ratio in the condenser is kept low on account of the high boiling point. Oil is stored in closed container preferable in Aluminum bottles.
- **Chemical Constituents:** The woody chips have an essential oil commonly known as Agar oil from 0.8% to 2.2% in fungal infested wood of 8-50 years old plant. The wood contains hexadecanoic acid (25.0%), pentadecanoic acid (6.7%) and oleic acid (4.9%); other constituents range from 0.1 to 2.1%.
- **Yield and Cost of Cultivation (Hectare):** This oil is exceptionally costly.

Therapeutic Uses

Wood is used as stimulant, aphrodisiac, tonic in diarrhea, vomiting and used in skin related ailments like wounds, injuries, pain, indigestion, heart related ailments, blood purifier against gout, impotence and urine related disorders. The plant acts as anti-inflammatory, stimulates the nervous system, antirheumatic and antiparalysis.

