

# *Psoralea corylifolia* Linn.

Syn. *Cullen corylifolium* (Linn.) Medik.

Fam. Fabaceae

<b>Ayurvedic name</b>	Bakuchi
<b>Unani name</b>	Babchi
<b>Hindi name</b>	Babchi, Bakuchi
<b>English name</b>	Purple Fleabane
<b>Trade name</b>	Bavchi, Bakuchi
<b>Parts used</b>	Seeds



*Psoralea corylifolia*

## Morphological Characteristics

**B**abchi is leguminous, erect, annual herb that grows 60-100 cm tall. The plant branches profusely and its stem and branches are covered with white hairs. Leaves are simple, 2.5-7.0 cm long, petiolate, rounded, with toothed margin and both sides covered with conspicuous black glandular dots. The seeds are surrounded by a sticky oily pericarp which contains psoralein.

## Floral Characteristics

Flowers are axillary, blue in colour and arranged in 10-30 flowered racemes. Calyx is nearly sessile. Corolla is yellowish-blue and little exerted. Pods are 5 cm long, subglobose, slightly compressed, closely pitted and beaked; seeds are oblong, flattened dark brown and covered with a mucilaginous layer. The seeds swell when placed in water. When the seeds are rubbed they give an aromatic odour and tastes slightly bitter.

## Distribution

It is mainly found in plains of Central India and Eastern part of Rajasthan, Punjab, and adjoining areas of Uttar Pradesh. It is sporadically cultivated in Rajasthan, Uttar Pradesh and Tamil Nadu States.



## Climate and Soil

The crops can be grown well in sub-tropical climate receiving low to medium rainfall over a variety of soils ranging from sandy medium loam to black cotton soils. Red loamy soil with good organic matters and a pH ranging from 6.5-7.5 are good for cultivation.

## Propagation Material

Seeds.

## Agro-technique<sup>26</sup>

### Nursery Technique

- **Raising Propagules:** No propagules are raised. The crop is raised through direct sowing of seeds which germinate easily. As a sole crop 8 kg seeds are needed for one hectare area.
- **Prapagule Rate and Pretreatment:** No specific pre-treatment is required for seeds before germination.

### Planting in the Field

- **Land Preparation and Fertilizer Application:** The land is prepared by ploughing 2-3 times with disc plough harrowed and planked to make the soil to a fine tilth before onset of monsoon. The area is divided into plots of convenient size. The main and sub-irrigation channels are laid out. NPK (Nitrogen, Phosphorous, Potassium) at the rate of 60: 60: 30 kg/ha are given as basal dose and mixed in the soil together with 10 t/ha of FYM.
- **Transplanting and Optimum Spacing:** Seeds are directly sown in lines at an optimum spacing of 60X30 cm to raise crop.
- **Intercropping System:** This crop can be cultivated as intercrop in tree plantation in orchards.
- **Interculture and Maintenance Practices:** Regular weeding (2-3) and hoeing operations are needed during early period of growth to control weeds.
- **Irrigation Practices:** The crop is rainfed and can stand partial drought conditions. However, 2-3 irrigations are required after sowing, depending on soil conditions and distribution of monsoon rains.
- **Disease and Pest Control:** Powder mildew is common problem during the winter months. Control measures involve spraying wetttable sulphur (sulfex) at the rate of 3% at weekly interval for 3 to 4 times. Leaf roller caterpillar is another menance and is controlled by 2-3 spray of 0.2% Endosulfan at fortnightly interval.

---

<sup>26</sup> Agro technique study carried out by Mahatma Phule Krishi Vidyapeeth (MPKV), Rahuri, Maharashtra.

